## **Injection Technique. The basics.**

#### **Preparing for injection**



Clean injection site and

(alcohol is not required)



Gather supplies.

Including your pen, insulin and pen needles.





Mix cloudy insulins: roll 10 times, then tip 10 times to see milky white consistency.







Attach your pen needle





Use pen needles and syringes only once.



Prime your pen.

Prime pen upwards with 2 or 3 units as per pen instructions. Repeat if needed until drops come out. (GLP-1 pens only need to be primed the 1st time you use them))

### Injecting







Choose injection site.

(injection into the back of your own arm is not recommended)

#### Remember









Rotate between injection areas

**Rotate within** injection zones







Dial dose. Insert, push completely and count to 10.





Insert the needle through the skin using a quick, smooth movement without excess force.



Withdraw the needle from the skin. Remove pen needle and dispose of it in an approved sharps container.





To ensure you are getting your insulin where it needs to go, to be properly absorbed, ask a health care professional to help determine the injection sites and technique that would be best for you.

### **Injection Technique. The basics.** Children Children (>6 years) & Children (>6 years) Adults Adults (2-6 years) Teens (slim) & Teens (slim) 4mm 5<sub>m</sub>m 6mm (pen or syringe) Preferable to use shorter pen needles and syringes 8mm 8mm (pen or syringe) 12.7mm Not Recommended

# Did you know?



There is a correct, and incorrect way to perform a skin lift. Delicately lift the skin and subcutaneous tissue, or fatty layer, between your thumb and index finger, leaving the muscle behind.





#### References

Gibney MA, et al. Skin and subcutaneous adipose layer thickness in adults with diabetes at sites used for insulin injections: implications for needle length recommendations. Curr Med Res Opin. 2010; 26 (6): 1519-1530.

 $\textbf{Berard L, et al.} \ \textbf{FIT Canada Recommendations for Injection Technique 4th Edition, September 2020}$ 

Lo Presti D, et al. Skin and subcutaneous thickness at injecting sites in children with diabetes: ultrasound findings and recommendations for giving injection. Ped Diab 2012.

Rini CR, et al. J Diabetes Sci Technol. 2019;13(3):533-545

